



**MEDICATION REFERENCE
NOT FOR USE DURING PATIENT CARE
Medications General**

Initial Date: 11/17/2023

Revised Date:

Section: MR23

Methylprednisolone

Pharmacological Category: Corticosteroid, Systemic

Routes: IV/IO/IM

Indications:

1. Allergic reactions
2. Airway inflammation
3. Reactive airway disease
4. Acute adrenal insufficiency

Contraindications:

1. Hypersensitivity to methylprednisolone (or similar)

Expected effects:

1. Decreased inflammation

Side effects:

1. Dizziness
2. Nausea/vomiting

Notes:

1. Prednisone PO is preferred over methylprednisolone for respiratory distress however prednisone it is not a required medication, and the PO tablet has restrictions (tablet cannot be cut, cannot be administered to children \leq 6 years of age, cannot be administered to patient that is unable to safely take PO medication).

Dosing: ANAPHYLAXIS ALLERGIC REACTION

Indication: If patient is symptomatic of an allergic reaction but not in a severe allergic reaction or anaphylaxis OR after epinephrine administration

Adults administer:

1. Methylprednisolone 125 mg IV/IO/IM

Pediatrics administer:

1. Methylprednisolone according to MI MEDIC cards.
2. If MI MEDIC cards are not available administer Methylprednisolone 2 mg/kg IV/IO/IM. Maximum dose 125 mg.

Dosing: ADRENAL CRISIS

Indication: Patients with a known history of adrenal insufficiency, experiencing signs of crisis.

Adults administer:

1. Methylprednisolone 125 mg IV/IO/IM

Pediatrics administer:

1. Methylprednisolone according to MI MEDIC cards.
2. If MI MEDIC cards are not available administer Methylprednisolone 2 mg/kg IV/IO/IM. Maximum dose 125 mg

REFERENCE ONLY – NOT FOR USE DURING PATIENT CARE



**MEDICATION REFERENCE
NOT FOR USE DURING PATIENT CARE
Medications General**

Initial Date: 11/17/2023

Revised Date:

Section: MR23

Dosing: ADULT RESPIRAOTRY DISTRESS

Indication: Respiratory distress patients with wheezing or diminished breath sounds due to asthma or COPD.

Adults administer:

1. Methylprednisolone 125 mg IV/IO/IM

Dosing: PEDIATRIC RESPIRATORY DISTRESS, FAILURE OR ARREST

Indication: Pediatric respiratory distress patients with suspected bronchospasm (wheezing)

Pediatrics administer:

1. Methylprednisolone according to MI MEDIC cards.
2. If MI MEDIC cards are not available administer Methylprednisolone 2 mg/kg IV/IO/IM.
Maximum dose 125 mg

Used in the Following Protocols:

Anaphylaxis/Allergic Reaction (Section 1 General Treatment)

Adrenal Crisis (Section 1 General Treatment)

Respiratory Distress (Section 3 Adult Treatment)

Pediatric Respiratory Distress, Failure or Arrest (Section 4 Obstetrics and Pediatrics)